



U.S.N

## P.E.S. College of Engineering, Mandya - 571 401

*(An Autonomous Institution affiliated to VTU, Belagavi)*

**Third Semester, B.E. - Semester End Examination; Dec - 2017/Jan - 2018**

### Engineering Mathematics - III

**(Common to all Branches)**

**Time: 3 hrs**

**Max. Marks: 100**

**Note:** Answer **FIVE** full questions, selecting **ONE** full question from each unit.

#### UNIT - I

- 1 a. Find the missing terms in the following table:

<i>x</i>	-0.2	0	0.2	0.4	0.6	0.8	1.0
<i>y</i>	2.6	-	3.4	4.28	-	14.2	29

6

- b. A survey conducted in a slum locality reveals the following information as classified below:

Income Per day (Rs.)	Under 10	10 - 20	20 - 30	30 - 40	40 - 50
Number of persons	20	45	115	210	115

7

Estimate the probable number of persons in the group 20 to 25.

- c. Fit an interpolating polynomial for the data using Newton's divided difference formula;

$u_0 = -5, u_1 = -14, u_4 = -125, u_8 = -21, u_{10} = 355$  and hence find  $u_2$ .

7

- 2 a. Find Lagrange's interpolating polynomial for the following data:

$$y(1) = 3, y(3) = 9, y(4) = 30, y(6) = 132.$$

6

- b. Given the data,

<i>x</i>	1	2	3	4	5
<i>y</i>	5	17	46	97	176

7

Find  $y(3.2)$ , by using Gauss forward formula.

- c. Apply Bessel's formula to find  $y_{25}$  given that  $y_{20} = 2854, y_{24} = 3162, y_{28} = 3544, y_{32} = 3922$ .

7

#### UNIT - II

- 3 a. Compute  $y''(1)$  for the function  $y = f(x)$  given:

<i>x</i>	0	2	4	6	8	10
<i>y</i>	0	4	56	204	496	980

6

- b. Find  $y'(3)$  from the following data by using Newton's divided difference formula,

<i>x</i>	0	1	2	5
<i>y</i>	2	3	12	147

7

- c. Find  $y''(2)$  by using Stirling's formula given,

<i>x</i>	-1	1	3	5
<i>y</i>	-3	21	77	165

7

- 4 a. Evaluate  $\int_0^1 \frac{dx}{1+x}$  by using Trapezoidal rule. Take seven ordinates.

6

- b. Use Simpson's  $\frac{1}{3}$  rd rule to find  $\int_0^{0.6} e^{-x^2} dx$  by taking six sub-intervals.

7

- c. Evaluate  $\int_{-\pi/2}^{\pi/2} \frac{dx}{2+\sin x}$  by using Weddle's rule by taking six equal parts.

7

**UNIT - III**

- 5 a. Obtain the Fourier series of the function  $f(x) = x(2\pi - x)$  in  $0 \leq x \leq 2\pi$ .

6

- b. Obtain the Fourier series of the function,

$$f(x) = x - x^2 \text{ in } (-\pi, \pi). \text{ Deduce that } \frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots$$

7

- c. Obtain the Fourier series of the function  $f(x) = |x|$  in  $(-l, l)$ . Hence deduce that,

$$\frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots$$

7

- 6 a. Obtain the complex Fourier series for the function  $f(x) = e^x$  in  $(0, 2l)$ .

6

- b. Obtain the cosine half-range series of  $f(x) = x \sin x$  in  $0 < x < \pi$ .

7

- c. Determine the constant term and first cosine and sine terms of the Fourier series expansion of the  $y$  from the following data:

$x^\circ$	$0^\circ$	$45^\circ$	$90^\circ$	$135^\circ$	$180^\circ$	$225^\circ$	$270^\circ$	$315^\circ$
$y$	2	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{1}{2}$

7

**UNIT - IV**

- 7 a. Find the Fourier transform of  $f(x) = xe^{-|x|}$ .

6

- b. Obtain the Fourier cosine transform of the function  $f(x) = \begin{cases} 4x, & 0 < x < 1 \\ 4-x, & 1 < x < 4 \\ 0, & x > 4 \end{cases}$ .

7

- c. Find the Fourier sine transform of  $f(x) = e^{-|x|}$  and hence evaluate  $\int_0^\infty \frac{x \sin mx}{1+x^2} dx$ ,  $m > 0$ .

7

- 8 a. Obtain the Z-transform of  $\cos n\theta$  and  $\sin n\theta$ .

6

- b. Obtain the inverse Z-transform of  $\frac{2z^2 + 3z}{(z+2)(z-4)}$ .

7

- c. Solve the difference equation  $y_{n+2} + 6y_{n+1} + 9y_n = 2^n$  with  $y_0 = 0, y_1 = 0$  using Z-transforms.

7

**UNIT - V**

- 9 a. Form the partial differential equation by eliminating the arbitrary constant from the relation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1.$$

6

- b. Solve  $x^2 \frac{\partial u}{\partial x} + y^2 \frac{\partial u}{\partial y} = 0$  by the method of separation of variables.

7

- c. Solve  $(mz - ny)p + (nx - lz)q = ly - mx$ .

7

- 10 a. Obtain the solution of the one dimensional heat equation  $\frac{\partial u}{\partial t} = c^2 \frac{\partial^2 u}{\partial x^2}$ .

10

- b. Solve the wave equation  $\frac{\partial^2 u}{\partial t^2} = c^2 \frac{\partial^2 u}{\partial x^2}$ , given that  $u(0, t) = 0, u(l, t) = 0, \frac{\partial u}{\partial t} = 0$

10

when  $t = 0$  and  $u(x, 0) = u_0 \sin\left(\frac{\pi x}{l}\right)$ .