



# P.E.S. College of Engineering, Mandya - 571 401

(An Autonomous Institution affiliated to VTU, Belagavi)

Second Semester, B.E. - Semester End Examination; May/June - 2018

## Engineering Mathematics - II

(Common to all Branches)

Time: 3 hrs

Max. Marks: 100

Note: Answer FIVE full questions, selecting ONE full question from each unit.

### UNIT - I

- 1 a. Investigate the values of  $\lambda$  and  $\mu$  such that the system of equations,  
 $x+y+z=6$ ,  $x+2y+3z=10$ ,  $x+2y+\lambda z=\mu$  has 6
- i) Unique solution      ii) Infinite many solution      iii) No solution
- b. Solve the following system of equations by using the LU - decomposition method: 7  
 $x+y+z=1$ ,  $3x+y-3z=5$  and  $x-2y-5z=10$ .
- c. Apply Gauss-Jordon method to solve  $x+2y+z=8$ ,  $2x+3y+4z=20$ , and  $4x+3y+2z=16$ . 7
- 2 a. State Cayley-Hamilton theorem and use it to compute the inverse of the matrix  $\begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 1 \\ 2 & 0 & 3 \end{bmatrix}$  6
- b. Find all the Eigen value and the Eigen vector corresponding to the least Eigen value of the matrix  $\begin{pmatrix} 6 & -2 & 2 \\ -2 & 3 & -1 \\ 2 & -1 & 3 \end{pmatrix}$  7
- c. Find the modal matrix P which diagonalizes the matrix  $A = \begin{bmatrix} 1 & 4 \\ 2 & 3 \end{bmatrix}$  7

### UNIT - II

- 3 a. Solve:  $\frac{d^2y}{dx^2} + 4\frac{dy}{dx} + 4y = 0$  given that  $y=0$ ,  $y'=-1$  at  $x=1$  6
- b. Solve:  $y'' + y' + y = x^2 + x + 1$ . 7
- c. Solve:  $\frac{d^2y}{dx^2} - 2\frac{dy}{dx} + 5y = e^{2x} \sin x$ . 7
- 4 a. Solve by the method of undetermined coefficients  $\frac{d^2y}{dx^2} + \frac{dy}{dx} - 2y = x + \sin x$ . 6
- b. Solve by the method of variation of Parameters  $\frac{d^2y}{dx^2} - 6\frac{dy}{dx} + 9y = \frac{e^{3x}}{x^2}$ . 7
- c. Solve  $(x+1)^2 \frac{d^2y}{dx^2} + (x+1) \frac{dy}{dx} + y = 4 \cos[\log(x+1)]$ . 7

### UNIT - III

- 5 a. Find the Laplace transform of (i)  $\frac{\sin at}{t}$  (ii)  $e^{3t} \sin 5t \sin 3t$ . 6

b. If  $f(t) = \begin{cases} t, & 0 \leq t \leq a \\ 2a-t, & a \leq t \leq 2a \end{cases}$ ,  $f(t+2a) = f(t)$ . Then show that  $L\{f(t)\} = \frac{1}{s^2} \tanh\left(\frac{as}{2}\right)$ . 7

c. Express  $f(t) = \begin{cases} \cos t, & 0 < t \leq \pi \\ 1, & \pi < t \leq 2\pi \\ \sin t, & t > 2\pi \end{cases}$  in terms of unit step function and hence find its Laplace 7

transform.

6 a. Find the inverse Laplace transform of (i)  $\frac{s+1}{s^2+s+1}$  (ii)  $\log\left(\frac{s+1}{s+2}\right)$ . 6

b. Find the inverse Laplace transform of  $\frac{s^2}{(s^2+a^2)(s^2+b^2)}$  by using convolution theorem. 7

c. Solve the Laplace transform method:  $\frac{d^2y}{dt^2} + 4\frac{dy}{dt} + 3y = e^{-t}$  with  $y(0) = 1$  and  $y'(0) = 1$ . 7

**UNIT - IV**

7 a. If  $x = r \cos \theta$  and  $y = r \sin \theta$  prove that  $JJ' = 1$ . 6

b. Expand  $e^x \log(1+y)$  in powers of  $x$  and  $y$  upto third degree terms by Taylor's theorem. 7

c. The temperature  $T$  at any point  $(x, y, z)$  in space is  $T = 400xyz^2$ . Find the highest temperature at the surface of the unit sphere  $x^2 + y^2 + z^2 = 1$ . 7

8 a. Evaluate  $\int \vec{F} \cdot d\vec{r}$  from  $(0, 0)$  to  $(1, 1)$  along (i)  $y = x$  (ii)  $y = \sqrt{x}$ , where  $\vec{F} = x^2i + xyj$ . 6

b. Apply Stoke's theorem to evaluate  $\oint_C (x+y)dx + (2x-z)dy + (y+z)dz$ , where  $C$  is the boundary of the triangle with vertices  $A = (2, 0, 0)$ ,  $B = (0, 3, 0)$  and  $C = (0, 0, 6)$ . 7

c. Apply Gauss divergence theorem to evaluate  $\iiint_s \vec{F} \cdot \hat{n} ds$  where  $\vec{F} = (x^2 - yz)\hat{i} + (y^2 - zx)\hat{j} + (z^2 - xy)\hat{k}$ . 7

Taken over the rectangular paralleopiped  $0 \leq x \leq a$ ,  $0 \leq y \leq b$ ,  $0 \leq z \leq c$ .

**UNIT - V**

9 a. Evaluate  $\int_0^4 \int_0^{\sqrt{z}} \int_0^{\sqrt{4z-x^2}} dy dx dz$ . 6

b. Evaluate  $\iint xy(x+y)dxdy$  taken over the area between  $y = x^2$  and  $y = x$ . 7

c. Evaluate  $\int_0^1 \int_0^{\sqrt{1-x^2}} (y^2)dxdy$  by changing the order of integration. 7

10 a. Find the area of the ellipse  $\frac{x^2}{9} + \frac{y^2}{16} = 1$  by double integration. 6

b. Evaluate the integral  $\iiint_R (xy)dxdydz$  where  $R$  is the positive octant of the sphere  $x^2 + y^2 + z^2 = a^2$ . 7

c. Define Beta function. Evaluate  $\int_0^4 x^{\frac{3}{2}}(4-x)^{\frac{5}{2}} dx$ . 7